

# (12) UK Patent Application (19) GB (11) 2 247 885 (13) A

(43) Date of A publication 18.03.1992

(21) Application No 9117175.1

(22) Date of filing 08.08.1991

(30) Priority data

(31) 9019933

(32) 12.09.1990

(33) GB

(71) Applicant

Farmitalia Carlo Erba S.r.l.

(Incorporated in Italy)

Via Carlo Imbonati 24, 20159 Milan, Italy

(72) Inventors

Alberto Bargiotti

Daniela Falardi

Antonino Suarato

(74) Agent and/or Address for Service

J A Kemp & Co

14 South Square, Gray's Inn, London, WC1R 5LX,  
United Kingdom

(51) INT CL<sup>5</sup>

C07H 19/02, A61K 31/71

(52) UK CL (Edition K)

C2C CAA CUA C1253 C1562 C1672 C215 C22X  
C22Y C25Y C253 C255 C28X C30Y C351 C353  
C36Y C360 C361 C362 C363 C364 C365 C388  
C50Y C509 C634 C635 C643 C644 C652 C662  
C672 C776 C80Y C801  
U1S S1313

(56) Documents cited

GB 2124224 A

EP 0128670 A

US 4672057 A

(58) Field of search


UK CL (Edition K) C2C CUA

INT CL<sup>5</sup> C07H

Online database : CAS ONLINE

(54) 13-dihydro-3'-(2-alkoxy-4-morpholinyl)anthracyclines

(57) Anthracycline glycosides of formula A:



wherein  $R_1$  is hydrogen atom, hydroxy or methoxy group;  $R_2$  or  $R_3$  represents hydroxyl group and the other of  $R_2$  and  $R_3$  represents hydrogen atom;  $R_4$  is hydrogen or hydroxy; both  $R_5$  and  $R_6$  represent hydrogen or one of  $R_5$  and  $R_6$  represents hydrogen and the other of  $R_5$  and  $R_6$  hydroxy;  $R_7$  represents a lower linear or branched alkyl residue or benzyl residue and their pharmaceutically acceptable salts are *anti-tumour agents*.

The invention relates to new anthracycline glycosides, to processes for their preparation and to 5 pharmaceutical compositions containing them.

wherein R<sub>1</sub> is a hydrogen atom, hydroxy or methoxy group; one  
10 of R<sub>2</sub> and R<sub>3</sub> represents a hydroxyl group and the other of R<sub>2</sub>  
and R<sub>3</sub> represents hydrogen; R<sub>4</sub> is hydrogen or hydroxy; both  
R<sub>5</sub> and R<sub>6</sub> represent hydrogen or one of R<sub>5</sub> and R<sub>6</sub> is hydroxy  
and the other of R<sub>5</sub> and R<sub>6</sub> is hydrogen; R<sub>7</sub> represents a  
lower linear or branched alkyl, preferably containing from 1  
15 to 10 carbon atoms, or benzyl group; or a pharmaceutically  
acceptable salt thereof.

Thus the compounds of the invention are 13-dihydro-anthracycline glycosides in which a 3'-nitrogen atom is

enclosed in a 2-alkoxy-4-morpholino ring.

The alkyl group represented by  $R_7$  may for example contain from 1 to 6, preferably 1 to 4, carbon atoms and may for instance be methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl or tert-butyl.

The compounds may be in the form of a mixture of isomers in which the 13-carbon atom has an S or an R configuration, such as the racemate.

Alternatively the compounds may be in optically pure form. The compounds may be in the S configuration at the 13-carbon atom and be substantially free of the isomer with the R configuration at the 13-carbon atom, or they may be in the R configuration at the 13-carbon atom and be substantially free of the isomer with the S configuration at the 13-carbon atom.


Particularly preferred compounds are those in which the 13-carbon atom has an S-configuration, i.e. wherein  $R_2 = OH$  and  $R_3 = H$ . Preferred pharmaceutically acceptable salts are acid addition salts such as hydrochlorides. Preferred embodiments of the new anthracycline glycosides of general formula A include:

A1: 13-(R/S)-dihydro-3'-deamino-3'-(2-methoxy-4-morpholinyl)-doxorubicin  
( $R_1 = OCH_3$ ,  $R_2 = OH/H$  and  $R_3 = H/OH$ ,  $R_4 = R_5 = OH$ ,  $R_6 = H$ ,  
25  $R_7 = CH_3$ )

A2: 13-(S)-dihydro-3'-deamino-3'-(2-methoxy-4-morpholinyl) doxorubicin

- A3: 13-(S/R)-dihydro-4'epi-3'-deamino-3'-(2-methoxy-4-morpholinyl)-doxorubicin  
( $R_1=OCH_3$ ,  $R_2=OH/H$  and  $R_3=H/OH$ ,  $R_4=R_6=OH$ ,  $R_5=H$ ,  $R_7=CH_3$ )
- 5 A4: 13-(S)-dihydro-4'epi-3'-deamino-3'-(2-methoxy-4-morpholinyl)-doxorubicin  
( $R_1=OCH_3$ ,  $R_2=OH$  and  $R_3=H$ ,  $R_4=R_6=OH$ ,  $R_5=H$ ,  $R_7=CH_3$ )
- A5: 13-(S/R)-dihydro-4-demethoxy-3'-deamino-3'-(2-methoxy-4-morpholinyl)daunorubicin  
10 ( $R_1=R_4=R_6=H$ ,  $R_2=OH/H$  and  $R_3=H/OH$ ,  $R_5=OH$ ,  $R_7=CH_3$ )
- A6: 13-(S)-dihydro-4-demethoxy-3'-deamino-3'-(2-methoxy-4-morpholinyl)daunorubicin  
( $R_1=R_4=R_6=H$ ,  $R_2=OH$  and  $R_3=H$ ,  $R_5=OH$ ,  $R_7=CH_3$ )

The dihydro-anthracyclines of the present invention  
15 may be prepared by several methods. The present invention, provides a first process for preparing a compound of formula A or a pharmaceutically acceptable salt thereof which first process comprises reducing the 13-carbonyl group of a compound of general formula B:



wherein R<sub>1</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub> and R<sub>7</sub> have the same meaning as above defined or a salt thereof, such as a pharmaceutically acceptable salt, eg the hydrochloride addition salt and, if desired, converting the resulting compound of formula A into  
5 a pharmaceutically acceptable salt.


The reduction may be effected using sodium borohydride in organic solvents such as methanol or using sodium cyanoborohydride for example in a mixture of acetonitrile and water typically at a pH from 7 to 4.

10 Preferably the reaction is carried out at 0°C for 5 minutes. This process affords a 1:1 mixture of 13(R)- and 13(S)- dihydro anthracyclines (Scheme 1). If the product is in the form of a free base it is preferably treated with methanolic hydrogen chloride and isolated as its hydrochloride. The  
15 starting compounds of formula B are described US-A-4,672,057 and our copending GB Patent Application No. 9007513.6, or in E.W. Acton in Bioactive Molecules, 55-101, vol 6, Edited by J.W. Lown, Elsevier 1988 and may be prepared by the methods described therein.

20 In particular GB Application No. 9007513.6 discloses compounds of formula B, in which R<sub>1</sub> is OMe, R<sub>4</sub> is OH, R<sub>5</sub> is OH, R<sub>6</sub> is H and R<sub>7</sub> is C<sub>2</sub>-C<sub>6</sub> alkyl and their pharmaceutically acceptable salts. It discloses that there may be prepared by

25 (i) reacting doxorubicin or an acid addition salt thereof, for example the hydrochloride salt, with a

diiodo compound of general formula C:




wherein R<sub>7</sub> is as defined above; and

(ii) if desired, converting the anthracycline glycoside of formula (A) thus obtained into a  
5 pharmaceutically acceptable acid addition salt thereof. It will be apparent to the person skilled in the art that analogous processes may be used to prepare other compounds of formula B and their salts.

The alkylation of the C-3' amino group of  
10 doxorubicin or the doxorubicin salt is typically performed in step (i) in a polar aprotic solvent and in the presence of a dry organic base such as triethylamine. Reaction is generally carried out at room temperature from eight to twenty four hours. The carbon atom C-2 bearing the -OR<sub>7</sub>  
15 group in the diiodo compound may have a (S) or (R) configuration. In a preferred embodiment, doxorubicin or its hydrochloride, dissolved in a polar aprotic solvent is reacted at room temperature and in the presence of a dry organic base, with the diiodo compound of general formula C  
20 to give the corresponding morpholinyl doxorubicin derivative of formula B which, after purification on a silica gel


column using as eluting system methylene chloride-methanol (97:5 v/v), is isolated, by treatment with methanolic anhydrous hydrogen chloride, as its hydrochloride.

The optically pure diiodo compounds C may be prepared starting from sugar precursors such as the compounds of general formula S derived from L-arabinose:




wherein  $R_7$  is as defined above. This process comprises:

(a) subjecting to periodate oxidation a compound of formula  $S^1$ :




10 wherein  $R_7$  is as defined above;

(b) reducing the thus-obtained dialdehyde derivative of formula  $T^1$ :



wherein  $R_7$  is as defined above;

(c) sulfonating the thus-obtained dihydroxy derivative of formula  $U^1$ :



wherein  $R_7$  is as defined above; and

5 (d) iodinating the sulfonated derivative thus obtained.


In order to prepare the diiodo compounds  $C$ , 1-substituted sugars  $S^1$ , prepared following standard procedures described in "Methods on Carbohydrate Chemistry" 10 Acad. Press., Vol 1, (1962), are first transformed into dialdehyde derivatives  $T^1$ . Generally, D- or L-arabinose is employed as a starting material. This is reacted with an alcohol  $R_7-OH$  thereby to form the compound of formula  $S^1$ . The dialdehyde derivatives can be obtained by using 15 periodate oxidation in water, then reduced to 1,5-dihydroxy-2-alkoxy or -benzyloxy-3-oxa-pentane  $U^1$  by using reducing agents such as sodium borohydride or sodium cyanoborohydride at pH 6.5 in a mixture of water and methanol.

The resultant dihydro compounds  $U^1$  are sulfonated 20 at the 1- and 5-hydroxyl groups, typically by using p-toluensulfonyl chloride in pyridine at 4°C to give the



sulfonyl ester of from which the diiodo derivatives C are obtained upon treatment with sodium or potassium iodide in aprotic solvent such as methylethylketone at 85°C from one to two days. The sequence of these reactions do not affect the chirality at C-2 of the diiodo derivatives C which is the same as the starting sugars S.


Other methods allow the preparation of optically pure 13(S)-dihydro anthracyclines of general formula A ( $R_2=OH$  and  $R_3=H$ ). Therefore the invention further provides a second process for the preparation of a compound of formula A, or a pharmaceutically acceptable salt thereof, which process comprises reacting an optically pure 13(S)-dihydro-anthracycline of general formula D:



D

15 wherein  $R_1$ ,  $R_4$ ,  $R_5$  and  $R_6$  have the same meaning as above defined,  $R_2$  represents hydroxyl group and  $R_3$  is hydrogen or a salt thereof, such as a pharmaceutically acceptable salt e.g. the hydrochloride addition salt with (a) diiodo or (b)

dialdehyde derivatives of general formula E:



E

wherein X represent iodine atom or formyl group (-CHO) and  
5  $R_7$  has the same meaning as above defined and, if desired,  
converting the resulting compound of formula A into to a  
pharmaceutically acceptable salt.

More particularly, compounds of general formula D  
may be alkylated (a) by using diiodo derivatives of formula  
10 E ( $X=I$ ) in the manner described in GB Patent Application No.  
9007513.6, in dry polar and aprotic solvents such as  
acetonitrile or dimethylformamide in the presence of a dry  
organic base, such as triethylamine typically at room  
temperature from 4 to 24 hours (Scheme 2), or may be  
15 reductively alkylated (b) using dialdehyde derivatives of  
formula E ( $X=CHO$ ) in aqueous media typically at pH from 5 to  
4 in the presence of a reducing agent such as sodium  
cyanoborohydride (Scheme 3).

If the product of either of these reactions is in  
20 the form of a free base it is preferably treated with  
methanolic hydrogen chloride and isolated as its  
hydrochloride.


The compounds of formula D may be prepared by as  
disclosed in US-A-4,438,105.

25 It will be appreciated that the compounds of  
formula E in which X is I correspond to the compounds of  
formula C described above and the compounds may be prepared


as described above. The compounds of formula E in which X is CHO may be prepared as disclosed in US-A-4,672,057 or by analogous methods which would be apparent to the person skilled in the art.

5 The following reaction Schemes illustrate the processes of the present invention.


Scheme 1



Scheme 2



Scheme 3



The present invention further provides a pharmaceutical composition comprising an anthracycline glycoside of formula A or a pharmaceutically acceptable salt thereof, such as the hydrochloride, together with a

5 pharmaceutically acceptable diluent or carrier. Such a composition may comprise conventional carriers and diluents and may be formulated and administered in conventional manner.

The compounds of the invention are useful in

10 methods of treatment of the human or animal body by therapy. They are useful as anti-tumour agents. A therapeutically effective amount is administered to a patient having a tumour to ameliorate or improve the condition of the patient. An amount sufficient to inhibit the growth of the

Biological Assay

13-(R/S)-dihydro-3'-deamino-3'[2(S)-methoxy-4-morpholinyl]doxorubicin (compound A1), was tested "in vitro" against LoVo and LoVo-resistant-doxorubicin (LoVo/DX) cells using a single cell plating technique after 4 hr treatment (Colony assay). The 50% inhibition concentration ( $IC_{50}$ ) was calculated on concentration-response curves. Compound A1 was tested in comparison with Doxorubicin and 3'-deamino-3'[2(S)-methoxy-4-morpholinyl]doxorubicin. Data are reported in Table 1.

Table 1: Cytotoxicity after 4 hr treatment  $IC_{50}$ =ng/ml<sup>(1)</sup>

Compound	LoVo	LoVo/DX	R.I. <sup>(2)</sup>
	$IC_{50}$ (ng/ml)	$IC_{50}$ (ng/ml)	
<u>A1</u>	28	118	4.2
Doxorubicin	60	2160	36
3'-deamino-3'- [2(S)-methoxy-4- morpholinyl] doxorubicin	16	33	2

<sup>(1)</sup>  $IC_{50}$  = concentration inhibiting 50% colony growth

<sup>(2)</sup> R.I. = Resistance Index =  $(IC_{50} \text{ LoVo/DX}) / (IC_{50} \text{ LoVo})$

Compound A1 was evaluated "in vivo" against P388 murine Leukemias, sensitive and resistant to Doxorubicin, in comparison with Doxorubicin and 3'-deamino-3'[2(S)-methoxy-4-morpholinyl]doxorubicin. Data are reported in Table 2.

Table 2: Antitumor activity against P388 and P388/DX  
(Johnson) Leukemias.

Compound	P388 <sup>(1)</sup>		P388/DX <sup>(2)</sup>	
	Dose <sup>(3)</sup> (mg/kg)	T/C <sup>(4)</sup> %	Dose <sup>(3)</sup> mg/kg	T/C <sup>(4)</sup> %
<u>A1</u>	1	161	1	133
Doxorubicin	13-16.9	200-225	13-16.9	86-100
3'-deamino-3'- [2(S)-methoxy-4- morpholinyl] doxorubicin	0.09	250	0.09	250

<sup>(1)</sup> 10<sup>6</sup> cells/mouse (P388 Leukemia) transplanted i.v. in  
CDF1 mice.

Treatment i.v. on day 1 after inoculation of tumor.

<sup>(2)</sup> 10<sup>5</sup> cells/mouse (P388/DX, Johnson) transplanted i.v. in  
CDF1 mice.

Treatment on day 1 after inoculation of tumor.

<sup>(3)</sup> Optimal Dose

<sup>(4)</sup> Median survival time; % over untreated controls.

5           The following Examples illustrate the invention.

Example 1

Preparation of 13-(R/S)dihydro-3'(2-methoxy-4-morpholinyl)  
doxorubicin (A1)

3'-deamino-3'(2-methoxy-4-morpholinyl)doxorubicin.HCl form  
10 (B1):  $R_1=OCH_3$ ,  $R_4=R_5=OH$ ,  $R_6=H$ ,  $R_7=CH_3$ ) (0.15 g, 0.22 mmole)  
was dissolved in methanol (25 ml), cooled at 0°C and treated  
with sodium borohydride (20 mg) under stirring. After five  
minutes, a mixture of acetone (10ml) and acetic acid (2ml)  
was added. The reaction mixture was diluted with water (50  
15 ml) and extracted twice with methylene chloride. After  
that, the aqueous solution was brought to pH 7.2 with  
aqueous hydrogen carbonate and extracted with methylene  
chloride. The organic phase was washed with water,  
separated, dried over anhydrous sodium sulphate, filtered  
20 and concentrated to small volume under reduced pressure.  
The title compound A1 (0.12g, yield 80%) was obtained by  
adding methanolic anhydrous hydrogen chloride followed by  
ethyl ether precipitation.

TLC on Kieselgel Plate F<sub>254</sub> (Merck), eluting system  
25 methylene chloride/methanol (6/1 by volume)  $R_f=0.52$

FD-MS: m/e 629 (M<sup>+</sup>)

<sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>);  $\delta$  1.38, 1.39 (d, J= 6.6 Hz, 3H, 5'-

CH<sub>3</sub>); 1.75 (m, 2H, 2'-CH<sub>2</sub>); 2.3-2.0 (m, 8H, CH<sub>2</sub>-N-CH<sub>2</sub>, 3'-H, 10ax-H, 8-CH<sub>2</sub>); 3.2-3.4 (m, 1H, 10e-H); 3.38 (s, 3H, O-CH-OCH<sub>3</sub>); 3.55 (m, 2H, NCH<sub>2</sub>-CH(H)O, 13-CH); 3.66 (m, 1H, 4'-H); 3.8-4.1 (m, 4H, 5'-H, CHOHCH<sub>2</sub>OH, NCH<sub>2</sub>CH(H)-O); 4.08 (s, 3H, 5 4-OCH<sub>3</sub>); 4.48 (m, 1H, O-CH-OCH<sub>3</sub>); 4.58, 4.63 (s, 1H, 9-OH); 5.28 (m, 1H, 7-H); 5.55 (1H, 1'3H); 7.38, (d, J=7.3Hz, 1H, 3-H); 7.78 (t, J=7.3Hz, 1H, 2-H); 8.03 (d, J=7.3Hz, 1H, 1-H); 13.32, 13.24 (s, 1H, 11-OH); 13.96, 13.97 (s, 1H, 6-OH).

Example 2

10 Preparation of 13-(S)dihydro-3'-deamino-3'(2-methoxy-4-morpholinyl)doxorubicin (A2)

13-(S)-dihydrodoxorubicin.HCl (C1: R<sub>1</sub>=OCH<sub>3</sub>, R<sub>2</sub>=OH and R<sub>3</sub>=H, R<sub>4</sub>=R<sub>5</sub>=OH, R<sub>6</sub>=H, R<sub>7</sub>=CH<sub>3</sub>) (0.10 g, 0.17 mmole) was dissolved in dry dimethylformamide (8 ml) and added with 1,5-diiodo-  
15 2(S)methoxyloxy-3-oxa-pentane (D1: X=I, R<sub>7</sub>=CH<sub>3</sub>) (0.5 g, 2 mmole) and dry triethylamine (0.5 ml, 0.4 mmole). The mixture was kept at room temperature for 24 hours, then was poured into water and extracted with methylene chloride. After standard work-up, the crude product was purified on  
20 silicic acid column using as eluting system a mixture of methylene chloride/ methanol (10/1 by volume), to give, after treatment with methanolic anhydrous hydrogen chloride, the title compound A2 (0.04 g, yield 40%).


TLC on Kieselgel Plate F<sub>254</sub> (Merck), eluting system  
25 methylene chloride/methanol (6/1 by volume) R<sub>f</sub>=0.52

FD-MS: m/e 629 (M<sup>+</sup>).



CLAIMS

1. An anthracycline glycoside of general formula A:



A

- 5 wherein  $R_1$  is a hydrogen atom, hydroxy or methoxy group; one of  $R_2$  and  $R_3$  represents a hydrogen group and the other of  $R_2$  and  $R_3$  represents a hydroxyl atom;  $R_4$  is hydrogen or hydroxy; both  $R_5$  and  $R_6$  represent hydrogen or one of  $R_5$  and  $R_6$  is hydroxy and the other of  $R_5$  and  $R_6$  represents
- 10 hydrogen;  $R_7$  represents a lower linear or branched alkyl or benzyl group or a pharmaceutically acceptable salt thereof.
2. A compound according to claim 1 in which  $R_7$  is an alkyl group containing from 1 to 6 carbon atoms.
3. A compound according to claim 2 which is
- 15 13-(R/S)-dihydro-3'-deamino-3'-(2-methoxy-4-morpholinyl)-doxorubicin or its hydrochloride salt.
4. A compound according to claim 2, which is 13-(S)-dihydro-3'-deamino-3'-(2-methoxy-4-morpholinyl)-

doxorubicin or its hydrochloride salt.


5. A compound according to claim 2, which is 13-(S/R)-dihydro-4'-epi-3'-deamino-3'-(2-methoxy-4-morpholinyl)-doxorubicin or its hydrochloride salt.

5 6. A compound according to claim 2, which is 13-(S)-dihydro-4'-epi-3'-deamino-3'-(2-methoxy-4-morpholinyl)-doxorubicin or its hydrochloride salt.

7. A compound according to claim 2, which is 13-(S/R)-dihydro-4-demethoxy-3'-deamino-3'-(2-methoxy-4-  
10 morpholinyl)-daunorubicin or its hydrochloride salt.

8. A compound according to claim 2, which is 13-(S)-dihydro-4-demethoxy-3'-deamino-3'-(2-methoxy-4-morpholinyl)-daunorubicin or its hydrochloride salt.

9. A process for preparing a compound  
15 according to claim 1, which process comprises reducing the 13-carbonyl group of a compound of formula B:




B

wherein R<sub>1</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub> and R<sub>7</sub> are as defined in claim 1, or

a salt thereof and, if desired, converting the resulting compound of formula A into a pharmaceutically acceptable salt.

10. A process according to claim 9, which comprises reacting a compound of formula A, with an alkaline metal borohydride or cyanoborohydride, in an organic solvent, at a temperature of 0°C for five minutes, to give the desired compound as a free base, treating with methanolic hydrogen chloride, and isolating the desired compound as a hydrochloride.

11. A process for preparing a compound according to claim 1, which process comprises reacting an optically pure 13(S)-dihydro anthracycline glycoside of formula D:




15

D

wherein R<sub>1</sub>, R<sub>4</sub>, R<sub>5</sub> and R<sub>6</sub> are as defined in claim 1, R<sub>2</sub> represents hydroxyl group and R<sub>3</sub> is hydrogen, or a salt

thereof, with a diiodo or dialdehyde of general formula E



E

wherein X represents an iodine atom or formyl group (-CHO)  
5 and R<sub>7</sub> is as defined in claim 1 and, if desired, converting  
the resulting product of formula A to a pharmaceutically  
acceptable salt.

12. A process according to claim 11, which is  
carried out: (a) if X represents iodine in a dry polar  
10 aprotic solvent and in the presence of a dry organic base at  
room temperature from 4 to 24 hours or (b) if X represents  
formyl, in an aqueous system, at pH from 5 to 4 in presence  
of a reducing agent and

in which if a free base is obtained it is treated  
15 with methanolic hydrogen chloride, to obtain a  
hydrochloride.

13. A pharmaceutical composition comprising a  
compound according to claim 1, together with a  
pharmaceutically acceptable diluent or carrier.

20 14. A compound as defined in claim 1 for use  
in a method of treatment of the human or animal body by  
therapy.

15. A compound according to claim 14 for use  
as an anti-tumour agent.

25 16. A process for the preparation of a  
compound as defined in claim 1, which is substantially as  
hereinbefore described in any one of the Examples.